arXiv:math/0103203v2 [math.AG] 8 Sep 2002

THE ENDOMORPHISM RINGS OF JACOBIANS OF CYCLIC COVERS OF THE PROJECTIVE LINE

BY YURI G. ZARHIN †

Department of Mathematics, Pennsylvania State University,
University Park, PA 16802,USA
e-mail: `zarhin@math.psu.edu`

Abstract. Suppose K is a field of characteristic zero, K_a is its algebraic closure, $f(x) \in K[x]$ is an irreducible polynomial of degree $n \geq 5$, whose Galois group coincides either with the full symmetric group $\mathbf{S}_n$ or with the alternating group $\mathbf{A}_n$. Let p be an odd prime, $\mathbf{Z}[\zeta_p]$ the ring of integers in the pth cyclotomic field $\mathbf{Q}(\zeta_p)$. Suppose C is the smooth projective model of the affine curve $y^p = f(x)$ and $J(C)$ is the jacobian of C. We prove that the ring $\mathrm{End}(J(C))$ of K_a-endomorphisms of $J(C)$ is canonically isomorphic to $\mathbf{Z}[\zeta_p]$.

1. Introduction

We write $\mathbf{Z}, \mathbf{Q}, \mathbf{C}$ for the ring of integers, the field of rational numbers and the field of complex numbers respectively. Recall that a number field is called a CM-field if it is a purely imaginary quadratic extension of a totally real field. Let p be an odd prime, $\zeta_p \in \mathbf{C}$ a primitive pth root of unity, $\mathbf{Q}(\zeta_p) \subset \mathbf{C}$ the pth cyclotomic field and $\mathbf{Z}[\zeta_p]$ the ring of integers in $\mathbf{Q}(\zeta_p)$. It is well-known that $\mathbf{Q}(\zeta_p)$ is a CM-field of degree $p-1$. We write $\mathbf{F}_p$ for the finite field consisting of p elements.

Let $f(x) \in \mathbf{C}[x]$ be a polynomial of degree $n \geq 4$ without multiple roots. Let $C_{f,p}$ be a smooth projective model of the smooth affine curve

$$y^p = f(x).$$

It is well-known that the genus $g(C_{f,p})$ of $C_{f,p}$ is $(p-1)(n-1)/2$ if p does not divide n and $(p-1)(n-2)/2$ if it does. The map

$$(x, y) \mapsto (x, \zeta_p y)$$

---

† Partially supported by the NSF.

gives rise to a non-trivial birational automorphism

$$\delta_p : C_{f,p} \to C_{f,p}$$

of period p.

The jacobian $J^{(f,p)} := J(C_{f,p})$ of $C_{f,p}$ is an abelian variety of dimension $g(C_{f,p})$. We write $\mathrm{End}(J^{(f,p)})$ for the ring of endomorphisms of $J^{(f,p)}$ over $\mathbf{C}$. By Albanese functoriality, δ_p induces an automorphism of $J^{(f,p)}$ which we still denote by δ_p; it is known ([10, p. 149], [11, p. 458]) that

$$\delta_p^{p-1} + \cdots + \delta_p + 1 = 0$$

in $\mathrm{End}(J^{(f,p)})$. This gives us an embedding

$$\mathbf{Z}[\zeta_p] \cong \mathbf{Z}[\delta_p] \subset \mathrm{End}(J^{(f,p)})$$

([10, p. 149], [11, p. 458]).

Our main result is the following statement.

Theorem 1.1. *Let K be a subfield of $\mathbf{C}$ such that all the coefficients of $f(x)$ lie in K. Assume also that $f(x)$ is an irreducible polynomial in $K[x]$ of degree $n \ge 5$ and its Galois group over K is either the symmetric group $\mathbf{S}_n$ or the alternating group $\mathbf{A}_n$. Then*

$$\mathrm{End}(J^{(f,p)}) = \mathbf{Z}[\delta_p] \cong \mathbf{Z}[\zeta_p].$$

In particular, $J^{(f,p)}$ is a simple complex abelian variety.

Remark 1.2. In the case when p is a Fermat prime the assertion of Theorem 1.1 is proven in [20]. (Also in [20] the author proved that if the conditions of Theorem 1.1 hold true then $\mathbf{Z}[\delta_p]$ is a maximal commutative subring in $\mathrm{End}(J^{(f,p)})$ for all odd primes p. See [21] for a similar result in positive characteristic when $p \mid n$ and $n \ge 9$.) The "opposite" case when $J^{(f,p)}$ is an abelian variety of CM-type was studied in [2]. An analogue of Theorem 1.1 for hyperelliptic jacobians (i.e., the case of $p = 2$) was proven in [17] (see also [18], [19]).

Examples 1.3.

1. the polynomial $x^n - x - 1 \in \mathbf{Q}[x]$ has Galois group $\mathbf{S}_n$ over $\mathbf{Q}$ ([13, p. 42]). Therefore the endomorphism ring (over $\mathbf{C}$) of the jacobian $J(C)$ of the curve $C : y^p = x^n - x - 1$ is $\mathbf{Z}[\zeta_p]$ if $n \ge 5$.
2. the Galois group of the "truncated exponential"

$$\exp_n(x) := 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \cdots + \frac{x^n}{n!} \in \mathbf{Q}[x]$$

is either $\mathbf{S}_n$ or $\mathbf{A}_n$ [12]. Therefore the endomorphism ring (over $\mathbf{C}$) of the jacobian $J(C)$ of the curve $C: y^p = \exp_n(x)$ is $\mathbf{Z}[\zeta_p]$ if $n \geq 5$.

Remark 1.4. If $f(x) \in K[x]$ then the curve $C_{f,p}$ and its jacobian $J^{(f,p)}$ are defined over K. Let $K_a \subset \mathbf{C}$ be the algebraic closure of K. Clearly, all endomorphisms of $J^{(f,p)}$ are defined over K_a. This implies that in order to prove Theorem 1.1, it suffices to check that $\mathbf{Z}[\delta_p]$ coincides with the ring of all K_a-endomorphisms of $J^{(f,p)}$ or equivalently, that $\mathbf{Q}[\delta_p]$ coincides with the $\mathbf{Q}$-algebra of K_a-endomorphisms of $J^{(f,p)}$.

The paper is organized as follows. Section 2 contains auxiliary results about endomorphism algebras of complex abelian varieties. We use them in Section 3 in order to study endomorphisms of $J^{(f,p)}$. In Section 4 we prove the main result. The short last Section contains corrigendum to [20].

The author would like to thank the referee for useful comments.

2. Complex abelian varieties

Throughout this section we assume that Z is a complex abelian variety of positive dimension. As usual, we write $\mathrm{End}^0(Z)$ for the semisimple finite-dimensional $\mathbf{Q}$-algebra $\mathrm{End}(Z) \otimes \mathbf{Q}$. We write $\mathfrak{C}_Z$ for the center of $\mathrm{End}^0(Z)$. It is well-known that $\mathfrak{C}_Z$ is a direct product of finitely many number fields. All the fields involved are either totally real number fields or CM-fields. Let $H_1(Z, \mathbf{Q})$ be the first rational homology group of Z; it is a $2\dim(Z)$-dimensional $\mathbf{Q}$-vector space. By functoriality $\mathrm{End}^0(Z)$ acts on $H_1(Z, \mathbf{Q})$; hence we have an embedding

$$\mathrm{End}^0(Z) \hookrightarrow \mathrm{End}_{\mathbf{Q}}(H_1(Z, \mathbf{Q}))$$

(which sends 1 to 1).

Suppose E is a subfield of $\mathrm{End}^0(Z)$ that contains the identity map. Then $H_1(Z, \mathbf{Q})$ becomes an E-vector space of dimension

$$d = \frac{2\dim(Z)}{[E : \mathbf{Q}]}.$$

We write

$$\mathrm{Tr}_E : \mathrm{End}_E(H_1(Z, \mathbf{Q})) \to E$$

for the corresponding trace map on the E-algebra of E-linear operators in $H_1(Z, \mathbf{Q})$.

Extending by $\mathbf{C}$-linearity the action of $\mathrm{End}^0(Z)$ and of E on the complex cohomology group

$$H_1(Z,\mathbf{Q})\otimes_{\mathbf{Q}}\mathbf{C}=H_1(Z,\mathbf{C})$$

of Z we get the embeddings

$$E\otimes_{\mathbf{Q}}\mathbf{C}\subset\mathrm{End}^0(Z)\otimes_{\mathbf{Q}}\mathbf{C}\hookrightarrow\mathrm{End}_{\mathbf{C}}(H_1(Z,\mathbf{C}))$$

which provide $H_1(Z,\mathbf{C})$ with a natural structure of free $E_{\mathbf{C}}:=E\otimes_{\mathbf{Q}}\mathbf{C}$-module of rank d. If Σ_E is the set of embeddings of $\sigma:E\hookrightarrow\mathbf{C}$ then it is well-known that

$$E_{\mathbf{C}}=E\otimes_{\mathbf{Q}}\mathbf{C}=\prod_{\sigma\in\Sigma_E}E\otimes_{E,\sigma}\mathbf{C}=\prod_{\sigma\in\Sigma_E}\mathbf{C}_\sigma$$

where

$$\mathbf{C}_\sigma=E\otimes_{E,\sigma}\mathbf{C}=\mathbf{C}.$$

Since $H_1(Z,\mathbf{C})$ is a free $E_{\mathbf{C}}$-module of rank d, there is the corresponding trace map

$$\mathrm{Tr}_{E_{\mathbf{C}}}:\mathrm{End}_{E_{\mathbf{C}}}(H_1(Z,\mathbf{C}))\to E_{\mathbf{C}}$$

which coincides on $E_{\mathbf{C}}$ with multiplication by d and with Tr_E on $\mathrm{End}_E(H_1(Z,\mathbf{Q}))$.

We write $\mathrm{Lie}(Z)$ for the tangent space of Z; it is a $\dim(Z)$-dimensional $\mathbf{C}$-vector space. By functoriality, $\mathrm{End}^0(Z)$ and therefore E acts on $\mathrm{Lie}(Z)$. This provides $\mathrm{Lie}(Z)$ with a natural structure of $E\otimes_{\mathbf{Q}}\mathbf{C}$-module. We have

$$\mathrm{Lie}(Z)=\bigoplus_{\sigma\in\Sigma_E}\mathbf{C}_\sigma\mathrm{Lie}(Z)=\oplus_{\sigma\in\Sigma_E}\mathrm{Lie}(Z)_\sigma$$

where

$$\mathrm{Lie}(Z)_\sigma=\mathbf{C}_\sigma\mathrm{Lie}(Z)=\{x\in\mathrm{Lie}(Z)\mid ex=\sigma(e)x\quad\forall e\in E\}.$$

Let us put

$$n_\sigma=n_\sigma(Z,E)=\dim_{\mathbf{C}_\sigma}\mathrm{Lie}(Z)_\sigma=\dim_{\mathbf{C}}\mathrm{Lie}(Z)_\sigma.$$

We write $\sigma':E\hookrightarrow\mathbf{C}$ for the composition of $\sigma:E\hookrightarrow\mathbf{C}$ and the complex conjugation $\mathbf{C}\to\mathbf{C}$. The embedding $\sigma'\in\Sigma_E$ is (called) the complex-conjugate of σ.

Remark 2.1. It is well-known ([1, p. 53], [5, p. 84]) that

$$n_\sigma+n_{\sigma'}=d\quad\forall\sigma\in\Sigma_E.$$

Remark 2.2. Let $\Omega^1(Z)$ be the space of the differentials of the first kind on Z. It is well-known that the natural map

$$\Omega^1(Z) \to \mathrm{Hom}_{\mathbf{C}}(\mathrm{Lie}(Z), \mathbf{C})$$

is an isomorphism. This isomorphism allows us to define via duality the natural homomorphism

$$E \to \mathrm{End}_{\mathbf{C}}(\mathrm{Hom}_{\mathbf{C}}(\mathrm{Lie}(Z), \mathbf{C})) = \mathrm{End}_{\mathbf{C}}(\Omega^1(Z)).$$

This provides $\Omega^1(Z)$ with a natural structure of $E \otimes_{\mathbf{Q}} \mathbf{C}$-module in such a way that

$$\Omega^1(Z)_\sigma := \mathbf{C}_\sigma \Omega^1(Z) \cong \mathrm{Hom}_{\mathbf{C}}(\mathrm{Lie}(Z)_\sigma, \mathbf{C}).$$

In particular,

$$n_\sigma = \dim_{\mathbf{C}}(\mathrm{Lie}(Z)_\sigma) = \dim_{\mathbf{C}}(\Omega^1(Z)_\sigma).$$

Theorem 2.3. *Suppose E contains $\mathfrak{C}_Z$. Then the tuple*

$$(n_\sigma)_{\sigma \in \Sigma_E} \in \prod_{\sigma \in \Sigma_E} \mathbf{C}_\sigma = E \otimes_{\mathbf{Q}} \mathbf{C}$$

lies in $\mathfrak{C}_Z \otimes_{\mathbf{Q}} \mathbf{C}$. In particular, if $E/\mathbf{Q}$ is Galois and $\mathfrak{C}_Z \ne E$ then there exists a nontrivial automorphism $\kappa : E \to E$ such that $n_\sigma = n_{\sigma\kappa}$ for all $\sigma \in \Sigma_E$.

Proof. The inclusion $\mathfrak{C}_Z \subset E$ implies that $\mathfrak{C}_Z$ is a field.

There is a canonical Hodge decomposition ([7, chapter 1], [1, pp. 52–53])

$$H_1(Z, \mathbf{C}) = H^{-1,0} \oplus H^{0,-1}$$

where $H^{-1,0}$ and $H^{0,-1}$ are mutually "complex conjugate" $\dim(Z)$-dimensional complex vector spaces. This splitting is $\mathrm{End}^0(Z)$-invariant and the $\mathrm{End}^0(Z)$-module $H^{-1,0}$ is canonically isomorphic to $\mathrm{Lie}(Z)$. Let

$$\mathfrak{f}_H : H_1(Z, \mathbf{C}) \to H_1(Z, \mathbf{C})$$

be the $\mathbf{C}$-linear operator in $H_1(Z, \mathbf{C})$ defined as follows.

$$\mathfrak{f}_H(x) = -x \quad \forall x \in H^{-1,0}; \quad \mathfrak{f}_H(x) = 0 \quad \forall x \in H^{0,-1}.$$

Clearly, $\mathfrak{f}_H$ commutes with $\mathrm{End}^0(Z)$ and therefore with E. Hence $\mathfrak{f}_H$ may be viewed as an endomorphism of the free $E_{\mathbf{C}}$-module $H_1(Z, \mathbf{C})$; clearly, its trace is the tuple

$$(-n_\sigma)_{\sigma \in \Sigma_E} \in \prod_{\sigma \in \Sigma_E} \mathbf{C}_\sigma = E_{\mathbf{C}}.$$

Suppose $MT = MT_Z \subset \mathrm{GL}_{\mathbf{Q}}(H_1(Z, \mathbf{Q})))$ is the Mumford-Tate group of (the rational Hodge structure $H_1(Z, \mathbf{Q})$ and of) Z ([1, 9, 16]). It is a connected reductive algebraic $\mathbf{Q}$-group that contains scalars and could be described as follows ([16, section 6.3]). Let $mt \subset \mathrm{End}_{\mathbf{Q}}(H_1(Z,\mathbf{Q})))$ be the $\mathbf{Q}$-Lie algebra of MT; it is a reductive algebraic linear $\mathbf{Q}$-Lie algebra which contains scalars and its natural faithful representation in $H_1(Z,\mathbf{Q})$ is completely reducible. In addition, mt is the *smallest* $\mathbf{Q}$-Lie subalgebra in $\mathrm{End}_{\mathbf{Q}}(H_1(Z,\mathbf{Q})))$ enjoying the following property: its complexification

$$mt_{\mathbf{C}} = mt \otimes_{\mathbf{Q}} \mathbf{C} \subset \mathrm{End}_{\mathbf{C}}(H_1(Z,\mathbf{C}))$$

contains scalars and $\mathfrak{f}_H$. It is well-known that the centralizer of MT (and therefore of mt) in $\mathrm{End}_{\mathbf{Q}}(H_1(Z,\mathbf{Q}))$ coincides with $\mathrm{End}^0(Z)$. This implies that the center $\mathfrak{c}$ of mt lies in $\mathfrak{C}_Z$. Since mt is reductive, it splits into a direct sum

$$mt = mt^{ss} \oplus \mathfrak{c}$$

of $\mathfrak{c}$ and a semisimple $\mathbf{Q}$-Lie algebra mt^{ss}. Clearly, mt lies in $\mathrm{End}_E(H_1(Z,\mathbf{Q}))$.

Since mt^{ss} is semisimple and the trace map Tr_E is a Lie algebra homomorphism, $\mathrm{Tr}_E(mt^{ss}) = \{0\}$. Since $\mathfrak{c} \subset \mathfrak{C}_Z \subset E$, we have $\mathrm{Tr}_E(\mathfrak{c}) \subset \mathfrak{C}_Z$ and therefore

$$\mathrm{Tr}_E(mt) \subset \mathfrak{C}_Z.$$

This implies easily that

$$\mathrm{Tr}_{E_{\mathbf{C}}}(mt_{\mathbf{C}}) \subset \mathfrak{C}_Z \otimes_{\mathbf{Q}} \mathbf{C}.$$

In particular, since $\mathfrak{f}_H \in mt_{\mathbf{C}}$, we have $\mathrm{Tr}_{E_{\mathbf{C}}}(\mathfrak{f}_H) \in \mathfrak{C}_Z \otimes_{\mathbf{Q}} \mathbf{C}$. But $\mathrm{Tr}_{E_{\mathbf{C}}}(\mathfrak{f}_H) = (-n_\sigma)_{\sigma\in\Sigma}$. This implies easily that

$$(n_\sigma)_{\sigma\in\Sigma_E} = -\mathrm{Tr}_{E_{\mathbf{C}}}(\mathfrak{f}_H) \in \mathfrak{C}_Z \otimes_{\mathbf{Q}} \mathbf{C}.$$

In order to prove the second assertion of the theorem, notice that its assumptions imply that $E/\mathfrak{C}_Z$ is a nontrivial Galois extension. If $\kappa : E \to E$ is a non-identity element of the Galois group $\mathrm{Gal}(E/\mathfrak{C}_Z)$ then one may easily check that

$$\mathfrak{C}_Z \otimes_{\mathbf{Q}} \mathbf{C} \subset \{(u)_{\sigma\in\Sigma_E} \in \prod_{\sigma\in\Sigma_E} \mathbf{C} = E_{\mathbf{C}} \mid u_\sigma = u_{\sigma\kappa} \quad \forall\sigma\}.$$

3. Cyclic covers and jacobians

Throughout this paper we fix an odd prime p and assume that K is a field of characteristic zero. We fix an algebraic closure K_a and write $\mathrm{Gal}(K)$ for the absolute Galois group $\mathrm{Aut}(K_a/K)$. We also fix in K_a a primitive pth root of unity ζ.

Let $f(x) \in K[x]$ be a separable polynomial of degree $n \geq 4$. We write $\mathfrak{R}_f$ for the set of its roots and denote by $L = L_f = K(\mathfrak{R}_f) \subset K_a$ the corresponding splitting field. As usual, the Galois group $\mathrm{Gal}(L/K)$ is called the Galois group of f and denoted by $\mathrm{Gal}(f)$. Clearly, $\mathrm{Gal}(f)$ permutes elements of $\mathfrak{R}_f$ and the natural map of $\mathrm{Gal}(f)$ into the group $\mathrm{Perm}(\mathfrak{R}_f)$ of all permutations of $\mathfrak{R}_f$ is an embedding. We will identify $\mathrm{Gal}(f)$ with its image and consider it as a permutation group of $\mathfrak{R}_f$. Clearly, $\mathrm{Gal}(f)$ is transitive if and only if f is irreducible in $K[x]$.

We refer the reader to [6, 20, 3, 8] for the definition and properties of the *heart* $(\mathbf{F}_p^{\mathfrak{R}_f})^{00}$ over the field $\mathbf{F}_p$ of the group $\mathrm{Gal}(f)$ acting on the set $\mathfrak{R}_f$. Here we just recall that $(\mathbf{F}_p^{\mathfrak{R}_f})^{00}$ is a finite-dimensional $\mathbf{F}_p$-vector space provided with a natural structure of $\mathrm{Gal}(f)$-module.

Let $C = C_{f,p}$ be the smooth projective model of the smooth affine K-curve

$$y^p = f(x).$$

So C is a smooth projective curve defined over K. The rational function $x \in K(C)$ defines a finite cover $\pi : C \to \mathbf{P}^1$ of degree p. Let $B' \subset C(K_a)$ be the set of ramification points. Clearly, the restriction of π to B' is an *injective* map $B' \hookrightarrow \mathbf{P}^1(K_a)$, whose image is the disjoint union of ∞ and $\mathfrak{R}_f$ if p does *not* divide $\deg(f)$ and just $\mathfrak{R}_f$ if it does. We write

$$B = \pi^{-1}(\mathfrak{R}_f) = \{(\alpha, 0) \mid \alpha \in \mathfrak{R}_f\} \subset B' \subset C(K_a).$$

Clearly, π is ramified at each point of B with ramification index p. We have $B' = B$ if and only if n is divisible by p. If n is not divisible by p then B' is the disjoint union of B and a single point $\infty' := \pi^{-1}(\infty)$. In addition, the ramification index of π at $\pi^{-1}(\infty)$ is also p. Using Hurwitz's formula, one may easily compute the genus $g = g(C) = g(C_{p,f})$ of C ([4, pp. 401–402], [14, proposition 1 on p. 3359], [10, p. 148]). Namely, g is $(p-1)(n-1)/2$ if p does *not* divide n and $(p-1)(n-2)/2$ if it does.

Remark 3.1. Assume that p does not divide n and consider the plane triangle (Newton polygon)

$$\Delta_{n,p} := \{(j,i) \mid 0 \leq j, \quad 0 \leq i, \quad pj + ni \leq np\}$$

with the vertices $(0,0)$, $(0,p)$ and $(n,0)$. Let $L_{n,p}$ be the set of integer points in the interior of $\Delta_{n,p}$. One may easily check that g coincides with the number of elements of $L_{n,p}$. It is also clear that for each $(j,i) \in L_{n,p}$

$$1 \le j \le n-1; \quad 1 \le i \le p-1; \quad p(j-1)+(j+1) \le n(p-i).$$

Elementary calculations ([4, theorem 3 on p. 403]) show that

$$\omega_{j,i} := x^{j-1}dx/y^{p-i} = x^{j-1}y^{i}dx/y^{p} = x^{j-1}y^{i-1}dx/y^{p-1}$$

is a differential of the first kind on C for each $(j,i) \in L_{n,p}$. This implies easily that the collection $\{\omega_{j,i}\}_{(j,i)\in L_{n,p}}$ is a basis in the space of differentials of the first kind on C.

There is a non-trivial birational K_a-automorphism of C

$$\delta_p : (x,y) \mapsto (x, \zeta y).$$

Clearly, δ_p^p is the identity map and the set of fixed points of δ_p coincides with B'.

Let $J^{(f,p)} = J(C) = J(C_{f,p})$ be the jacobian of C. It is a g-dimensional abelian variety defined over K and one may view (via Albanese functoriality) δ_p as an element of

$$\mathrm{Aut}(C) \subset \mathrm{Aut}(J(C)) \subset \mathrm{End}(J(C))$$

such that $\delta_p \ne \mathrm{Id}$ but $\delta_p^p = \mathrm{Id}$ where Id is the identity endomorphism of $J(C)$. Here $\mathrm{Aut}(C)$ stands for the group of K_a-automorphisms of C, $\mathrm{Aut}(J(C))$ stands for the group of K_a-automorphisms of $J(C)$ and $\mathrm{End}(J(C))$ stands for the ring of all K_a-endomorphisms of $J(C)$. As usual, we write $\mathrm{End}^0(J(C)) = \mathrm{End}^0(J^{(f,p)})$ for the corresponding $\mathbf{Q}$-algebra $\mathrm{End}(J(C)) \otimes \mathbf{Q}$.

Lemma 3.2. $\mathrm{Id} + \delta_p + \cdots + \delta_p^{p-1} = 0$ *in* $\mathrm{End}(J(C))$. *Therefore the subring* $\mathbf{Z}[\delta_p] \subset \mathrm{End}(J(C))$ *is isomorphic to the ring* $\mathbf{Z}[\zeta_p]$ *of integers in the pth cyclotomic field* $\mathbf{Q}(\zeta_p)$. *The* $\mathbf{Q}$*-subalgebra* $\mathbf{Q}[\delta_p] \subset \mathrm{End}^0(J(C)) = \mathrm{End}^0(J^{(f,p)})$ *is isomorphic to* $\mathbf{Q}(\zeta_p)$.

Proof. See [10, p. 149], [11, p. 458].

Remark 3.3. If K contains ζ then the Galois modules $(\mathbf{F}_p^{\mathfrak{R}_f})^{00}$ and $\ker(\mathrm{Id} - \delta_p)$ are canonically isomorphic ([10, proposition 6.2], [11, proposition 3.2]).

Remark 3.4. Recall that p is odd and assume that $n = \deg(f)$ is divisible by p say, $n = pm$ for some positive integer m. Since $n \ge 4$, we conclude that $n \ge 5$.

Let $\alpha \in K_a$ be a root of f and $K_1 = K(\alpha)$ be the corresponding subfield of K_a. We have $f(x) = (x-\alpha)f_1(x)$ with $f_1(x) \in K_1[x]$. Clearly, $f_1(x)$ is a separable polynomial over K_1 of degree $pm - 1 = n - 1 \geq 4$. It is also clear that the polynomials

$$h(x) = f_1(x + \alpha), h_1(x) = x^{n-1}h(1/x) \in K_1[x]$$

are separable of the same degree $pm - 1 = n - 1 \geq 4$. The standard substitution

$$x_1 = 1/(x - \alpha), y_1 = y/(x - \alpha)^m$$

establishes a birational isomorphism between $C_{f,p}$ and a curve

$$C_{h_1} : y_1^p = h_1(x_1)$$

(see [14, p. 3359]). But $\deg(h_1) = pm - 1$ is *not* divisible by p. Clearly, this isomorphism commutes with the actions of δ_p.

Theorem 3.5. *Suppose* $n \geq 4$. *Assume that* $\mathbf{Q}[\delta_p]$ *is a maximal commutative subalgebra in* $\mathrm{End}^0(J^{(f,p)})$. *Then the center* $\mathfrak{C}$ *of* $\mathrm{End}^0(J^{(f,p)})$ *is a CM-subfield of* $\mathbf{Q}[\delta_p]$.

Proof. This is theorem 3.8 of [20].

Theorem 3.6. *Suppose* $n \geq 4$. *Assume that* $\mathbf{Q}[\delta_p]$ *is a maximal commutative subalgebra in* $\mathrm{End}^0(J^{(f,p)})$. *Then* $\mathrm{End}^0(J^{(f,p)}) = \mathbf{Q}[\delta_p] \cong \mathbf{Q}(\zeta_p)$ *and therefore* $\mathrm{End}(J^{(f,p)}) = \mathbf{Z}[\delta_p] \cong \mathbf{Z}[\zeta_p]$.

Proof. Let $\mathfrak{C} = \mathfrak{C}_{J(f,p)}$ be the center of $\mathrm{End}^0(J^{(f,p)})$. We know that $\mathfrak{C}$ is a CM-subfield of $E := \mathbf{Q}[\delta_p]$.

Replacing, if necessary, K by its subfield (finitely) generated over $\mathbf{Q}$ by all the coefficients of f, we may assume that K (and therefore K_a) is isomorphic to a subfield of the field $\mathbf{C}$ of complex numbers. So, $K \subset K_a \subset \mathbf{C}$. We may also assume that $\zeta = \zeta_p$ and consider $C_{(f,p)}$ as complex projective curve and its jacobian $J^{(f,p)}$ as complex abelian variety.

Let $\Sigma = \Sigma_E$ be the set of all field embeddings $\sigma : E = \mathbf{Q}[\delta_p] \hookrightarrow \mathbf{C}$. We are going to apply Theorem 2.3 to $Z = J^{(f,p)}$ and $E = \mathbf{Q}[\delta_p]$. In order to do that we need to get some information about the multiplicities

$$n_\sigma = n_\sigma(Z, E) = n_\sigma(J^{(f,p)}, \mathbf{Q}[\delta_p]).$$

Remark 2.2 allows us to do it, using the action of $\mathbf{Q}[\delta_p]$ on the space $\Omega^1(J^{(f,p)})$ of differentials of the first kind on $J^{(f,p)}$.

Recall that if $\sigma' : \mathbf{Q}[\delta_p] \hookrightarrow \mathbf{C}$ is the embedding complex conjugate to σ then, by Remark 2.1,

$$n_\sigma + n_{\sigma'} = \frac{2\mathrm{dim}(J^{(f,p)})}{p-1},$$

since $[\mathbf{Q}[\delta_p] : \mathbf{Q}] = p - 1$. Notice also that for each $\sigma : \mathbf{Q}[\delta_p] \hookrightarrow \mathbf{C}$

$$\Omega^1(J^{(f,p)})_\sigma = \{\omega \in \Omega^1(J^{(f,p)}) \mid \delta_p(\omega) = \sigma(\delta_p)\omega\}.$$

In other words, $\Omega^1(J^{(f,p)})_\sigma$ is the eigenspace corresponding to the eigenvalue $\sigma(\delta_p)$ of δ_p and n_σ is the multiplicity of the eigenvalue $\sigma(\delta_p)$.

Let $i < p$ be a positive integer and $\sigma_i : \mathbf{Q}[\delta_p] \hookrightarrow \mathbf{C}$ be the embedding which sends δ_p to ζ^{-i}. Obviously, the complex conjugate of σ_i coincides with σ_{p-i}. In addition, for each σ there exists precisely one i such that $\sigma = \sigma_i$. Clearly, $\Omega^1(J^{(f,p)})_{\sigma_i}$ is the eigenspace of $\Omega^1(J^{(f,p)})$ attached to the eigenvalue ζ^{-i} of δ_p. Therefore n_{σ_i} coincides with the multiplicity of the eigenvalue ζ^{-i}.

Let P_0 be one of the δ_p-invariant points (i.e., a ramification point for π) of $C_{f,p}(K_a) \subset C_{f,p}(\mathbf{C})$. Then

$$\tau : C_{f,p} \to J^{(f,p)}, \quad P \mapsto \mathrm{cl}((P) - (P_0))$$

is an embedding of complex algebraic varieties and it is well-known that the induced map

$$\tau^* : \Omega^1(J^{(f,p)}) \to \Omega^1(C_{f,p})$$

is a $\mathbf{C}$-linear isomorphism obviously commuting with the actions of δ_p. (Here cl stands for the linear equivalence class.) This implies that n_{σ_i} coincides with the dimension of the eigenspace of $\Omega^1(C_{(f,p)})$ attached to the eigenvalue ζ^{-i} of δ_p.

Remark 3.7. Clearly, if for some positive integer j the differential $x^{j-1}dx/y^{p-i}$ lies in $\Omega^1(C_{(f,p)})$ then it is an eigenvector of δ_p with eigenvalue ζ^i. Now assume that p does *not* divide n. It follows from Remark 3.1 that each $n_\sigma = n_{\sigma_i}$ could be visualized as the number of interior integer points in $\Delta_{n,p}$ along the corresponding (to $p - i$) horizontal line. Elementary calculations show that this number is $[\frac{ni}{p}]$. This implies that $n_{\sigma_i} = [\frac{ni}{p}]$ for $1 \le i \le p-1$. Then $n_{\sigma_i} = 0$ if and only if $1 \le i \le [\frac{p}{n}]$.

Assume, in addition, that $p < n$. Clearly, in this case the function $i \mapsto n_{\sigma_i} = [\frac{ni}{p}]$ is strictly increasing.

Remark 3.8. Assume that p divides n. Then $n \ge 5$ and $n - 1 \ge 4$. Clearly, p does *not* divide $n - 1$. Applying Remark 3.4, we get a curve $C_{h_1,p} : y_1^p = h_1(x_1)$

with separable polynomial $h_1(x_1)$ of degree $n-1$ and a δ_p-equivariant birational isomorphism between $C_{f,p}$ and $C_{h_1,p}$. This gives us a δ_p-equivariant isomorphism

$$\Omega^1(C_{f,p}) \cong \Omega^1(C_{h_1,p}).$$

Applying Remark 3.7 to $C_{h_1,p}$ and $n-1$ (instead of $C_{f,p}$ and n), we conclude that $n_{\sigma_i} = [\frac{(n-1)i}{p}]$ for $1 \le i \le p-1$. Then $n_{\sigma_i} = 0$ if and only if $1 \le i \le [\frac{p}{n-1}]$.

Assume, in addition, that $n \ne p$. Clearly, in this case $n-1 > p$ and the function $i \mapsto n_{\sigma_i} = [\frac{(n-1)i}{p}]$ is strictly increasing.

Proposition 3.9. (i) *let us assume that $p > n$. Then $n_\sigma = 0$ if and only if $\sigma = \sigma_i$ with $1 \le i \le [\frac{(p-1)}{n}]$.*

(ii) *let us assume that $p = n$. Then $n_\sigma = 0$ if and only if $\sigma = \sigma_i$ with $i = 1$.*

Proof of Proposition 3.9. First, assume that $p > n$. Clearly, p does *not* divide n and therefore $[\frac{p}{n}] = [\frac{p-1}{n}]$. Now the assertion (i) follows from Remark 3.7.

Now assume that $n = p$. By Remark 3.8, $n_{\sigma_i} = 0$ if and only if $1 \le i \le [\frac{p}{n-1}]$. But $[\frac{p}{n-1}] = [\frac{p}{p-1}] = 1$.

Proposition 3.10. *Let us assume that $p < n$. If σ, ι are two embeddings $\mathbf{Q}[\delta_p] \hookrightarrow \mathbf{C}$ then $n_\sigma = n_\iota$ if and only if $\sigma = \iota$.*

Proof of Proposition 3.10. First assume that p does *not* divide n. Then the assertion follows from (the last sentence of) Remark 3.7.

Now assume that p divides n. Then the assertion follows from (the last sentence of) Remark 3.8.

End of the proof of Theorem 3.6. If $\mathfrak{C} = \mathbf{Q}[\delta_p]$ then we are done, since $\mathbf{Q}[\delta_p]$ is a maximal commutative subalgebra in $\mathrm{End}^0(J^{(f,p)})$. Assume that $\mathfrak{C} \ne \mathbf{Q}[\delta_p]$. Our goal is to get a contradiction.

Clearly, $\mathbf{Q}[\delta_p]/\mathbf{Q}$ is a Galois extension. It follows from Theorem 2.3 and Remark 2.2 (applied to $Z = J^{(f,p)}$ and $E = \mathbf{Q}[\delta_p]$) that there exists a non-trivial field automorphism $\kappa : \mathbf{Q}[\delta_p] \to \mathbf{Q}[\delta_p]$ such that for all $\sigma \in \Sigma$

$$n_\sigma = n_{\sigma\kappa}.$$

Clearly, there exists an integer m such that $1 < m < p$ and $\kappa(\delta_p) = \delta_p^m$.

First, assume that $n > p$. It follows from Proposition 3.10 that $\sigma\kappa = \sigma$ which could not be the case, since κ is *not* the identity map. This contradiction proves the Theorem in the case of $n > p$.

Second, assume that $n = p$. It follows from Proposition 3.9(ii) that $\sigma_1 \kappa = \sigma_1$ which, by the same token, leads to a contradiction.

Third, assume that $p > n$. It follows from Proposition 3.9(i) that the map $\sigma \mapsto \sigma\kappa$ permutes the set $\{\sigma_i \mid 1 \le i \le [\frac{(p-1)}{n}]\}$. Since $\kappa(\delta_p) = \delta_p^m$, $\sigma_i\kappa(\delta_p) = \zeta^{-im}$. This implies that multiplication by m in $\mathbf{F}_p^*$ leaves invariant the subset $A := \{i \bmod p \in \mathbf{F}_p \mid 1 \le i \le [\frac{(p-1)}{n}]\}$. This implies that

$$m = m \cdot 1 \le \left[\frac{(p-1)}{n}\right] \le \frac{(p-1)}{4}.$$

Let us consider the arithmetic progression consisting of the m integers $[\frac{(p-1)}{n}] + 1, \dots, [\frac{(p-1)}{n}] + m$ with difference 1. All its elements lie between $[\frac{(p-1)}{n}] + 1$ and

$$\left[\frac{(p-1)}{n}\right] + m \le 2\left[\frac{(p-1)}{n}\right] \le 2\frac{(p-1)}{4} = \frac{(p-1)}{2} < p - 1.$$

Clearly, there exists a positive integer $r \le m$ such that $[\frac{(p-1)}{n}] + r$ is divisible by m, i.e., there is a positive integer d such that $md = [\frac{(p-1)}{n}] + r$. Since $[\frac{(p-1)}{n}] \ge m \ge 2$, we have $d \le [\frac{(p-1)}{n}]$ but $md = [\frac{(p-1)}{n}] + r \le [\frac{(p-1)}{n}] + m < p - 1$. This implies that A is *not* invariant under multiplication by m which gives the desired contradiction.

4. Jacobians and their endomorphism rings

Recall that K is a field of characteristic zero, K_a is its algebraic closure. Suppose $f(x) \in K[x]$ is a polynomial of degree $n \ge 5$ without multiple roots, $\mathfrak{R}_f \subset K_a$ is the set of its roots, $K(\mathfrak{R}_f)$ is its splitting field. Let us put

$$\mathrm{Gal}(f) = \mathrm{Gal}(K(\mathfrak{R}_f)/K) \subset \mathrm{Perm}(\mathfrak{R}_f).$$

Theorem 4.1 (corollary 5.3 of [20]). *Let p be an odd prime. If $f(x) \in K[x]$ is an irreducible polynomial of degree $n \ge 5$ and $\mathrm{Gal}(f) = \mathbf{S}_n$ or $\mathbf{A}_n$ then $\mathbf{Q}[\delta_p]$ is a maximal commutative subalgebra in $\mathrm{End}^0(J^{(f,p)})$ and the center of $\mathrm{End}^0(J^{(f,p)})$ is a CM-subfield of $\mathbf{Q}[\delta_p]$.*

Combining Theorems 4.1 and 3.6, we obtain the following statement.

Theorem 4.2. *Let p be an odd prime. If $f(x) \in K[x]$ is an irreducible polynomial of degree $n \ge 5$ and $\mathrm{Gal}(f) = \mathbf{S}_n$ or $\mathbf{A}_n$ then $\mathrm{End}^0(J^{(f,p)}) = \mathbf{Q}[\delta_p]$ and therefore $\mathrm{End}(J^{(f,p)}) = \mathbf{Z}[\delta_p] \cong \mathbf{Z}[\zeta_p]$.*

Clearly, Theorem 1.1 is a special case of Theorem 4.2.

Example 4.3. Suppose $L = \mathbf{C}(z_1, \cdots, z_n)$ is the field of rational functions in n independent variables $z_1, \cdots, z_n$ with constant field $\mathbf{C}$ and $K = L^{\mathbf{S}_n}$ is the subfield of symmetric functions. Then $K_a = L_a$ and

$$f(x) = \prod_{i=1}^{n} (x - z_i) \in K[x]$$

is an irreducible polynomial over K with Galois group $\mathbf{S}_n$. Let C be a smooth projective model of the K-curve $y^p = f(x)$ and $J(C)$ its jacobian. It follows from Theorem 4.2 that if $n \ge 5$ then the ring of L_a-endomorphisms of $J(C)$ is $\mathbf{Z}[\zeta_p]$. In particular, the abelian variety $J(C)$ is absolutely simple. When $p = 3$ and $3 \mid n$ the absolute simplicity of $J(C)$ was proven in ([15, p. 107]).

Example 4.4. Let $h(x) \in \mathbf{C}[x]$ be a *Morse polynomial* of degree $n \ge 5$. This means that the derivative $h'(x)$ of $h(x)$ has $n-1$ distinct roots $\beta_1, \cdots \beta_{n-1}$ and $h(\beta_i) \ne h(\beta_j)$ while $i \ne j$. (For example, $x^n - x$ is a Morse polynomial.) Let $K = \mathbf{C}(z)$ be the field of rational functions in variable z with constant field $\mathbf{C}$ and K_a its algebraic closure. Then a theorem of Hilbert ([13, theorem 4.4.5, p. 41]) asserts that the Galois group of $h(x) - z$ over $k(z)$ is $\mathbf{S}_n$. Let C be a smooth projective model of the K-curve $y^p = h(x) - z$ and $J(C)$ its jacobian. It follows from Theorem 4.2 that the ring of K_a-endomorphisms of $J(C)$ is $\mathbf{Z}[\zeta_p]$. In particular, the abelian variety $J(C)$ is absolutely simple.

We refer the reader to [18, 19, 20, 22] for the definition and basic properties of *very simple* representations.

Theorem 4.5. *Suppose p is an odd prime, $n \ge 5$ and K contains a primitive pth root of unity. If the* $\mathrm{Gal}(f)$*-module* $(\mathbf{F}_p^{\mathfrak{R}_f})^{00}$ *is very simple then* $\mathbf{Q}[\delta_p]$ *coincides with its own centralizer in* $\mathrm{End}^0(J^{(f,p)})$.

Proof. See theorem 5.2 of [20].

Theorem 4.6. *Suppose p is an odd prime, $n \ge 5$ and K contains a primitive pth root of unity. If the* $\mathrm{Gal}(f)$*-module* $(\mathbf{F}_p^{\mathfrak{R}_f})^{00}$ *is very simple then* $\mathrm{End}^0(J^{(f,p)}) = \mathbf{Q}[\delta_p]$ *and therefore* $\mathrm{End}(J^{(f,p)}) = \mathbf{Z}[\delta_p] \cong \mathbf{Z}[\zeta_p]$.

Proof. It is an immediate corollary of Theorem 4.5 combined with Theorem 3.6.

5. CORRIGENDUM TO [20]

Remark 2.1 on p. 94, the last assertion. In general, it is not necessarily true that G is doubly transitive [3, Beispiel 2c], [8]. However, it becomes true if one assumes additionally that either p does not divide n or G is transitive and p is an odd number dividing n ([3, Satz 4a and Satz 11], [20, lemma 2.4]).

Lemma 2.4 on p. 95. Its assertion is essentially contained in Satz 4a of [3].

Remark 2.5 on p. 95. Its assertion is essentially Hilffsatz 3b of [3].

Sections 1, 3 and 5. Everywhere $\mathbf{Q}(\delta_p)$ means $\mathbf{Q}[\delta_p]$. (However, it does not make a difference, since $\mathbf{Q}[\delta_p]$ is a field.)

References

[1] P. Deligne, Hodge cycles on abelian varieties (notes by J.S. Milne). Lecture Notes in Math., vol. **900** (Springer-Verlag, 1982), pp. 9–100.

[2] J. de Jong and R. Noot, Jacobians with complex multiplications. In: *Arithmetic algebraic geometry* (eds. G. van der Geer, F. Oort and J. Steenbrink). Progress in Math., vol. **89** (Birkhäuser, 1991), pp. 177–192.

[3] M. Klemm, Über die Reduktion von Permutationsmoduln. *Math. Z.* **143** (1975), 113–117.

[4] J. K. Koo, On holomorphic differentials of some algebraic function field of one variable over C. *Bull. Austral. Math. Soc.* **43** (1991), 399–405.

[5] B. Moonen and Yu. G. Zarhin, Weil classes on abelian varieties. *J. reine angew. Math.* **496** (1998), 83–92.

[6] B. Mortimer, The modular permutation representations of the known doubly transitive groups. *Proc. London Math. Soc.* (3) **41** (1980), 1–20.

[7] D. Mumford, *Abelian varieties*, 2nd edn (Oxford University Press, 1974).

[8] P. M. Neumann, Review of [3], *MR* 52 #544.

[9] K. Ribet, Hodge classes on certain abelian varieties. *Amer. J. Math.* **105** (1983), 523–538.

[10] B. Poonen and E. Schaefer, Explicit descent for Jacobians of cyclic covers of the projective line. *J. reine angew. Math.* **488** (1997), 141–188.

[11] E. Schaefer, Computing a Selmer group of a Jacobian using functions on the curve. *Math. Ann.* **310** (1998), 447–471.

[12] I. Schur, Gleichungen ohne Affect. *Sitz. Preuss. Akad. Wiss.* 1930, *Physik-Math. Klasse* 443–449 (=Ges. Abh. III, 191–197).

[13] J.-P. Serre, *Topics in Galois Theory* (Jones and Bartlett Publishers, 1992).

[14] C. Towse, Weierstrass points on cyclic covers of the projective line. *Trans. Amer. Math. Soc.* **348** (1996), 3355-3377.

[15] H. Völklein, Cyclic covers of $\mathbf{P}^1$ and Galois actions on their division points. *Contemporary Math.* **186** (1994), 91–107.

[16] Yu. G. Zarhin, Weights of simple Lie algebras in the cohomology of algebraic varieties. *Izv. Akad. Nauk SSSR Ser. Mat.* **48** (1984), 264–304; English translation: *Math. USSR Izv.* **24** (1985), 245 - 281.

[17] Yu. G. Zarhin, Hyperelliptic jacobians without complex multiplication. *Math. Res. Letters* **7** (2000), 123–132.

[18] Yu. G. Zarhin, Hyperelliptic jacobians and modular representations. In: *Moduli of abelian varieties* (eds. C. Faber, G. van der Geer and F. Oort). Progress in Math., vol. **195** (Birkhäuser, 2001), pp. 473–490.

[19] Yu. G. Zarhin, Hyperelliptic jacobians without complex multiplication in positive characteristic. *Math. Res. Letters* **8** (2001), 429–435.

[20] Yu. G. Zarhin, Cyclic covers of the projective line, their jacobians and endomorphisms. *J. reine angew. Math.* **544** (2002), 91–110.

[21] Yu. G. Zarhin, Endomorphism rings of certain jacobians in finite characteristic. *Matem. Sbornik* **193** (2002), issue 8 (Russian), to appear.

[22] Yu. G. Zarhin, Very simple 2-adic representations and hyperelliptic jacobians; available at http://arXiv.org/abs/math.AG/0109014; *Moscow Math. J.* **2** (2002), issue 2, to appear.